Rotation About a Tilted Axis. Pensacola, Fla, Naval Aerospace Medical Research Laboratory - 1129, 1971

- 54. Denise P: Rotation d'axe incline par rapport a la gravite—Effets des petits angles et interet clinique, Thesis. Paris, Universite Pierre et Marie Curie, 1986
- 55. Kamerer DB, Wall C III, Furman JMR: Earth Horizontal Axis Rotational Responses in Patients With Unilateral Peripheral Vestibular Deficits (Abstr). Barany Society Meeting, Bologna, Italy, 1987
- 56. Wilson VJ, Jones GM: Mammalian Vestibular Physiology. New York, Plenum Press, 1979
- 57. Wall C III, Black FO: Postural stability and rotational tests: Their effectiveness for screening dizzy patients. Acta Otolaryngol 1983; 95:235-246
- 58. Bles W, de Jong JMBV: Uni- and bilateral loss of vestibular function, In Bles W, Brandt TH (Eds): Disorders of Posture and Gait. Amsterdam, Elsevier, 1986, pp 127-139
- $59.\;$  Nashner LM: A model describing vestibular detection of body sway motion. Acta Otolaryngol 1971; 72:429-436
- 60. Nashner LM, Black FO, Wall C III: Adaptation to altered support and visual conditions during stance: Patients with vestibular deficits. J Neurosci 1982; 2:536-544

- 61. Black FO, Wall C III, Nashner LM: Effects of visual and support surface orientation references upon postural control in vestibular deficit subjects. Acta Otolaryngol (Stockh) 1983; 95:199-201
- 62. Elidan J, Sohmer H, Nizan M: Recording of short latency vestibular evoked potentials to acceleration in rats by means of skin electrodes. Electroencephalogr Clin Neurophysiol 1982; 53:501-505
- 63. Elidan J, Langhofer L, Honrubia V: Recording of short-latency vestibular evoked potentials induced by acceleration impulses in experimental animals: Current status of the method and its application. Electroencephalogr Clin Neurophysiol 1987; 68:58-69
- 64. Salamy J, Potvin A, Jones K, et al: Cortical evoked responses to labyrinthine stimulation in man. Psychophysiology 1975; 12:55-61
- 65. Hofferberth B: Evoked potentials to rotatory stimulation. Acta Otolaryngol 1984; 406 (suppl): 134-136
- 66. Hood JD, Kayan A: Observations upon the evoked responses to natural vestibular stimulation. Electroencephalogr Clin Neurophysiol 1985; 62:266-276
- 67. Pirodda E, Ghedini S, Zanetti MA: Investigations into vestibular evoked responses. Acta Otolaryngol (Stockh) 1987; 104:77-84

## Travelers' Diarrhea

If you take a group of travelers from the United States to Mexico, you can show that a prophylactic antibiotic such as trimethoprim-sulfamethoxazole will prevent gastroenteritis in the treated group versus the placebo group. I don't think there's any doubt about that. (Studies show that Pepto-Bismol does the same thing.) The problem with making the recommendation generically and having all travelers carrying around a 7-day, a 10-day, or 2-week supply of trimethoprim-sulfamethoxazole is that it will lead to resistance and lessen the use of a very valuable drug combination. Most gram-negative rods are sensitive to trimethoprim-sulfamethoxazole, including the *E coli* in your own GI tract, and you may run into a *Salmonella* or a *Shigella* which is resistant to the trimethoprim-sulfamethoxazole. So now you've ingested a salad that has a *Shigella sonnei* or a *Shigella dysenteriae*, worse yet. And you are taking an antibiotic that is wiping out your own normal flora, which does have some counterbalancing effect in your fight against invading pathogens. And not only that, the antibiotic that you're taking is not effective against the organism that you've run into because it is resistant to the trimethoprim-sulfamethoxazole. This is the one that bothers me most, personally, when I'm tempted to take Bactrim if I'm going to Mexico.

-JOHN E. CONTE, Jr, MD

Extracted from Audio-Digest Internal Medicine, Vol. 35, No. 18, in the Audio-Digest Foundation's series of tape-recorded programs. For subscription information: 1577 E Chevy Chase Dr, Glendale, CA 91206